

Refractory Diarrhoea in Hyaline Fibromatosis Syndrome: A 5-Month-Old Infant Stabilized Through Precision Gut-Directed Nutritional Therapy

Chaitra^{1*}, Babu Ratnakar²

¹ICU Medical Officer, Department of Pediatrics, Government General Hospital (GGH), Khammam, Telangana, India

²Head of Department, Department of Pediatrics, Government General Hospital (GGH), Khammam

ABSTRACT

Hyaline Fibromatosis Syndrome (HFS) is an uncommon autosomal recessive disorder characterized by progressive deposition of hyaline material across multiple organ systems. In infants, the presentation may be subtle initially, but progression often leads to significant gastrointestinal complications. This case highlights a 5-month-old infant presenting with severe refractory diarrhoea, profound feeding difficulty, failure to thrive, and *E. coli* urosepsis. Conventional management failed, prompting the introduction of a structured gut-directed nutritional therapy involving semi-elemental feeds, osmotic load reduction, and carefully titrated fluid-electrolyte management. The infant demonstrated marked improvement, emphasizing the importance of aligning nutritional strategies with the underlying mucosal pathology in HFS.

Keywords: Hyaline Fibromatosis Syndrome; Refractory Diarrhoea; Gut-Directed Nutritional Therapy; Semi-Elemental Feeding; Infant Nutrition; Gastrointestinal Manifestations

INTRODUCTION

Hyaline Fibromatosis Syndrome (HFS) encompasses a spectrum of disorders caused by pathogenic variants in *ANTXR2*, resulting in the deposition of amorphous eosinophilic hyaline material in the skin, gastrointestinal tract, and musculoskeletal system. Although dermatological and joint-related manifestations often dominate the clinical picture, gastrointestinal symptoms may lead to the most severe complications, including dehydration, electrolyte imbalance, malnutrition, and recurrent infections [1]. The pathophysiological basis involves disruption of the lamina propria, reduced absorptive surface, mucosal stiffness, and impaired intestinal motility. Despite these significant implications, formal guidelines for the management of diarrhoea in HFS remain limited. This case sheds light on a physiology-driven nutritional intervention that may improve outcomes in similar clinical scenarios [2].

CASE PRESENTATION

A 5-month-old male infant, born at term with no perinatal complications, demonstrated progressive symptoms beginning at approximately two months of age. The earliest findings included

irritability during handling, firm plaques over the scalp and neck, and the appearance of small papulonodular lesions near the perianal and auricular areas [3]. These manifestations gradually worsened and were accompanied by feeding difficulty and faltering growth, raising concerns for an underlying systemic disorder. At 3½ months of age, dermatology evaluation noted prominent indurated plaques and rubbery nodules. A biopsy of these lesions revealed deposition of eosinophilic hyaline material, confirming the diagnosis of Hyaline Fibromatosis Syndrome [4]. At the time of acute admission, the child was severely dehydrated, lethargic, and markedly underweight. He had experienced persistent watery diarrhoea for ten days, leading to oliguria and worsening irritability. Laboratory investigations demonstrated leukocytosis, elevated inflammatory markers, metabolic acidosis, hyponatremia, and urine culture positive for *E. coli*. Standard management was initiated, including intravenous antibiotics, fluid correction using balanced crystalloids, zinc supplementation, probiotics, and lactose-free feeding. However, the diarrhoea did not improve and feed tolerance remained poor. Repeat assessments revealed progressive electrolyte derangements, raising concern for gastrointestinal mucosal dysfunction due to hyaline deposition.

***Correspondence to:** Chaitra, ¹ICU Medical Officer, Department of Pediatrics, Government General Hospital (GGH), Khammam, Telangana, India, Email: chaitrasri07@gmail.com

Received: May 22, 2026; **Manuscript No:** JPNB-26-3426; **Editor Assigned:** May 26, 2026; **PreQc No:** JPNB-26-3426 (PQ); **Reviewed:** May 28, 2026; **Revised:** September 01, 2026; **Manuscript No:** JPNB-26-3426 (R); **Published:** September 11, 2026

Citation: Chaitra, Ratnakar B (2026) Refractory Diarrhoea in Hyaline Fibromatosis Syndrome: A 5-Month-Old Infant Stabilized Through Precision Gut-Directed Nutritional Therapy. *J. Pediatr. Neonatal Sci.* Vol.2 Iss.2, September (2026), pp:67-68.

Copyright: © 2026 Chaitra, Babu Ratnakar. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[5] Given this pathophysiological background, a tailored gut-directed nutritional strategy was implemented [6]. This included a transition to semi-elemental formula to reduce digestive burden, small-volume frequent feeds to optimize absorption, strict osmotic reduction, and precise electrolyte replacement. Within 48–72 hours, the child demonstrated substantial clinical improvement, including decreased stool frequency, improved hydration, better feed tolerance, and normalization of urine output. He was subsequently discharged with a structured feeding plan and regular outpatient follow-up [7].

DISCUSSION

The gastrointestinal system is particularly vulnerable in HFS due to the accumulation of hyaline material within the mucosal layers. This disrupts nutrient absorption, weakens mucosal integrity, and impairs peristaltic movement. These structural changes explain the persistent and refractory nature of diarrhoea seen in affected infants, often rendering conventional paediatric diarrhoea protocols ineffective. In this case, standard management modalities showed minimal impact, emphasizing the unique nature of diarrhoea in HFS. The notable improvement following the introduction of semi-elemental feeds supports the hypothesis that reducing the digestive and antigenic load may substantially improve gastrointestinal handling. Additionally, meticulous fluid and electrolyte management proved essential, given the susceptibility of these patients to rapid deterioration. This case underscores the need for heightened clinical suspicion of HFS in infants presenting with early cutaneous lesions, joint stiffness, and unexplained diarrhoea. It also demonstrates that targeted nutritional strategies may significantly alter the trajectory of disease progression and reduce the risk of severe complications such as sepsis.

CONCLUSION

This case reinforces the importance of individualized nutritional strategies in managing gastrointestinal manifestations of Hyaline Fibromatosis Syndrome. Precision gut-directed therapy, involving semi-elemental feeding and electrolyte-optimized hydration, may offer a practical and effective intervention for infants with refractory diarrhoea. Early recognition and prompt management can mitigate complications and improve clinical outcomes in this vulnerable population.

ETHICAL APPROVAL AND PATIENT CONSENT

Written informed consent was obtained from the patient's parent/legal guardian for publication of this case report and accompanying clinical information/images. Patient identity has

been adequately anonymized in accordance with ethical publishing standards and GlobalMeet journal guidelines.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

FUNDING

No external funding was received for this study.

REFERENCES

1. Denadai R, Raposo-Amaral CE, Bertola D, Kim C, Alonso N, Hart T, Han S, Stelini RF, Buzzo CL, Raposo-Amaral CA, Hart PS. Identification of 2 novel ANTXR2 mutations in patients with hyaline fibromatosis syndrome and proposal of a modified grading system. *American journal of medical genetics Part A*. 2012 Apr; 158(4):732-42.
2. Mantri MD, Pradeep MM, Kalpesh PO, Pranavsinh RJ. Hyaline Fibromatosis Syndrome: A Rare Inherited Disorder. *Indian J Dermatol*. 2016 Sep-Oct;61(5):580. doi: 10.4103/0019-5154.190129. PMID: 27688461; PMCID: PMC5029258.
3. Raeekarami SR, Aghighi Y, Afshin A, Malek A, Zamani A, Ziaee V. Infantile Systemic Hyalinosis: Report of 17-year Experience. *Iran J Pediatr*. 2014 Dec;24(6):775-8. Epub 2014 Dec 9. PMID: 26019786; PMCID: PMC4442842.
4. Prathiba Rajalakshmi P, Srinivasan K. Gastrointestinal manifestations of Henoch-Schonlein purpura: A report of two cases. *World J Radiol*. 2015 Mar 28;7(3):66-9. doi: 10.4329/wjr.v7.i3.66. PMID: 25825636; PMCID: PMC4374091.
5. Rodrigues-Fernandes CI, de Cáceres CB, Sant'Ana MS, Soares CD, de Carvalho MG, van Heerden WF, Robinson L, Radhakrishnan R, Hunter KD, Gomez RS, de Almeida OP. Oral lesions containing amyloid-like material. *Oral surgery, oral medicine, oral pathology and oral radiology*. 2021 Aug 1;132(2):190-201.
6. Braegger C, Decsi T, Dias JA, Hartman C, Kolacek S, Koletzko B, Koletzko S, Mihatsch W, Moreno L, Puntis J, Shamir R, Szajewska H, Turck D, van Goudoever J; ESPGHAN Committee on Nutrition. Practical approach to paediatric enteral nutrition: a comment by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr*. 2010 Jul;51(1):110-22. doi: 10.1097/MPG.0b013e3181d336d2. PMID: 20453670.
7. Troeger CE, Khalil IA, Blacker BF, Biehl MH, Albertson SB, Zimsen SR, Rao PC, Abate D, Ahmadi A, Ibrahim Ahmed ML, Akal CG. Quantifying risks and interventions that have affected the burden of diarrhoea among children younger than 5 years: an analysis of the Global Burden of Disease Study 2017. *The Lancet Infectious Diseases*. 2020 Jan 1;20(1):37-59.